



Rx FOR SUCCESS

Clots and Emboli

Deep vein thromboses (DVT) are blood clots that form in the larger veins deep within the lower extremities or the pelvic area. A blood clot in a superficial vein (often in a varicose vein) has little mortality risk but can lead to local soft tissue inflammation (known as thrombophlebitis), ulceration, and infection (known as cellulitis). A blood clot in a deep vein may break off in pieces that travel first to the heart and then to the lungs. In the lungs, this event is called a pulmonary embolism (PE). A large clot to the lungs can cause acute pulmonary hypertension, low oxygen, and sudden death. Smaller clots can cause symptoms but may not cause sudden death. The death rate in recurrent PE is 45%.

DVT often causes leg pain and swelling, while the classic symptoms of PE are sudden onset of chest pain, shortness of breath, and cough productive of bloody sputum. Diagnosis is difficult and dangerous episodes can be misdiagnosed. Possible tests to detect DVT and PE include MRI, CT scanning, blood tests, ventilation-perfusion scan, venogram, and compression ultrasound of legs. Pulmonary angiogram is the best test for PE.

Risk factors for DVT (and subsequent PE) are stasis, trauma, and hypercoagulability. Stasis is associated with events that cause decreased mobility and/or reduced blood flow (such as being bedridden or in a leg cast). Soft tissue injury may cause direct damage to the vein. Any history, however old, of lower body trauma or surgery puts a person at risk for DVT. Hypercoagulable states are discussed under bleeding and clotting disorders (See Hypercoagulable Clotting Disorders).

Initial treatment for clots and emboli is usually hospitalization and anticoagulation therapy with injections of Heparin or Heparin-like products such as Lovenox and Fragmin. This is followed by oral anticoagulant with Coumadin for six months. In some cases, lifelong Coumadin treatment is required. Recurrent episodes are sometimes surgically treated with an intravascular filter (a Greenfield filter) to catch clots in the blood stream before they reach the lungs.

The increased mortality associated with clots is due to complications. These are infection, pulmonary embolism, and potentially dangerous bleeding episodes caused by the anticoagulant therapy.

UNDERWRITING CONSIDERATIONS

DVT or PE present	Postpone until resolved
Single episode of DVT or PE, resolved, on no medication	0
PE with current anticoagulant therapy (e.g., Coumadin, Heparin*)	Table B
Recurrent PE	
Two episodes in past 3 years	Table B
More than two episodes in last 3 years	Individual consideration
More than 3 years ago	Generally not rated
Varicose veins of legs, superficial thrombophlebitis	Generally not rated unless complicated by ulceration or infection

* Anyone on Coumadin, if not rated, is excluded from Preferred class. Those on Heparin will be rated a minimum of Table B.

This material is designed to provide general information about the subject matter covered. It should be used with the understanding that we are not rendering legal, accounting, or tax advice. Such services should be provided by the client's professional advisors. Accordingly, any information in this document cannot be used by any taxpayer for purposes of avoiding penalties under the Internal Revenue Code.

This material is intended for insurance informational purposes only and is not personal medical advice for clients. Rates and availability will vary based on the satisfaction of our underwriting criteria. Underwriting rules are subject to change at our discretion. This marketing material is subject to an expiration date, and use of this material must be discontinued as of the expiration date.

Insurance issued by The Prudential Insurance Company of America and its affiliates, Newark, NJ.

NOT FOR CONSUMER USE.

© 2013 Prudential Financial, Inc. and its related entities.
0191000-00002-00 Ed. 01/2013 Exp. 01/31/2015 Rx 127

Securities and Insurance Products:

Not Insured by FDIC or Any Federal Government Agency.
May Lose Value.

Not a Deposit of or Guaranteed by Any Bank or Bank Affiliate.



Prudential
Bring Your Challenges[®]

Ask “Rx”pert Underwriter (Ask Our Expert)

After reading the *Rx for Success* on Clots and Emboli, use this form to Ask “Rx”pert Underwriter for an informal quote.

Producer _____ Phone _____ Fax _____
 Client _____ Age/DOB _____ Sex _____

If your client has a history of clots or emboli, please answer the following:

1. Please list date of diagnosis.

2. Please note type of treatment.

- Coumadin Hospitalization _____ (Date)
 Aspirin Heparin

3. Was there a thromboembolic event?

- MI DVT Other
 CVA PE None

4. Has there been any evidence of recurrence?

- Yes. Please give details. _____
 No

5. Is your client on any other medications?

- Yes. Please give details. _____
 No

6. Has your client smoked cigarettes in the last 12 months?

- Yes. Please give details. _____
 No

7. Does your client have any other major health problems (e.g., cancer, etc.)?

- Yes. Please give details. _____
 No