

## **Depression and Anxiety Disorders**

Mood and anxiety disorders are common, and the mortality risk is due primarily to suicide, cardiovascular disease, and substance abuse. Risk is highest early in the course of the disorder or within 2 years of a hospitalization.

Mood disorders are divided into **Depressive Disorders** (*unipolar*) and **Bipolar Disorders** (*manic depressive*). **Dysthymia** is chronic low-grade depression that does not meet the criteria for **Major Depression**. Criteria for **Major Depression** require a history of depressed mood for at least 2 weeks plus 4 or more of the following: weight change, sleep disturbance, psychomotor agitation or retardation, fatigue, feelings of worthlessness or guilt, difficulty concentrating, or suicidal ideation. To meet the criteria for **Bipolar Disorder**, there must be a history of at least one episode of mania (*abnormal elevated/irritable mood*) in addition to the **Major Depression** criteria.

Anxiety disorders include panic disorders, agoraphobia, social phobia, social anxiety disorder (SAD), simple phobia, generalized anxiety disorder (GAD), obsessive-compulsive disorder (OCD), and post-traumatic stress disorder (PTSD). Symptoms include worry and nervousness, racing heart, breathlessness, dizziness, sweats, headache, insomnia, and other vague complaints. Depressive disorders often overlap with anxiety disorders, and in the long term, many patients continue to have symptoms. Recurrences are common for both mood and anxiety disorders.

Drug therapy (with or without counseling) is effective in treating most individuals. The mainstay of therapy for both anxiety and mood disorders is antidepressant drugs. For more severe cases, electroconvulsive therapy (ECT) or anti-psychotic agents may be required. If anxiety is present, treatment may include benzodiazepines and buspirone. Benzodiazepines are addicting and underwriting caution is necessary, especially in those prone to substance abuse. Bipolar disorder (manic depression) is usually treated with Lithium.

This material is designed to provide general information about the subject matter covered. It should be used with the understanding that we are not rendering legal, accounting, or tax advice. Such services should be provided by the client's professional advisors. Accordingly, any information in this document cannot be used by any taxpayer for purposes of avoiding penalties under the Internal Revenue Code.

This material is intended for insurance informational purposes only and is not personal medical advice for clients. Rates and availability will vary based on the satisfaction of our underwriting criteria. Underwriting rules are subject to change at our discretion. This marketing material is subject to an expiration date, and use of this material must be discontinued as of the expiration date.

Insurance issued by The Prudential Insurance Company of America and its affiliates, Newark, NJ.

NOT FOR CONSUMER USE.

© 2013 Prudential Financial, Inc. and its related entities. 0192751-00002-00 Ed. 01/2013 Exp. 01/15/2015 Rx 026

## **Securities and Insurance Products:**

Not Insured by FDIC or Any Federal Government Agency. May Lose Value.

Not a Deposit of or Guaranteed by Any Bank or Bank Affiliate.



UNDERWRITING CONSIDERATIONS FOR ADULTS WITH MOOD AND ANXIETY DISORDERS			
Mild	No suicidal ideation for 1yr, able to perform normal activities with minimal symptoms on $0-2$ medication (no antipsychotic agent), rare anxiety attacks, no ECT for >19yr, no hospitalization/suicide attempts/disability for >9yr	Non-rated	
Moderate	No suicidal ideation for 6 months, able to perform normal activities with minimal-moderate symptoms on 0 – 3 medications (no antipsychotic agent), single episode of disability (of short duration), no hospitalization/ECT/suicide attempts for >9yr	Table B	
Severe	Significant symptoms, requires antipsychotic medication, ECT/hospitalization(s) long episode(s) of disability, single suicide attempt	<ul> <li>Declined within first yr of recovery</li> <li>2nd – 3rd yr—Table D – F</li> <li>4th – 5th yr—Table B – D</li> <li>6th – 9th yr—Table B – C</li> <li>Higer rating for those with multiple severe episodes</li> </ul>	

Applicants under age 18, with a history of drug or alcohol abuse, with psychotic conditions, with multiple suicide attempts, or with frequent panic attacks will be given individual consideration.

To get an idea of how a client with Mood and Anxiety Disorders would be viewed in the underwriting process, use the Ask "Rx" pert Underwriter on the next page for an informal quote.

Ask "Rx"pert Underwriter (Ask Our Expe	ert)			
After reading the Rx for Success on Mood and	Anxiety Disorders, use this f	form to Ask "Rx" pert Underwriter for an informal quote.		
Producer	Phone	Fax		
Client	Age/DOB	Sex		
If your client has a history of mood or anxiety disorder, please answer the following:				
1. Please provide the diagnosis.				
2. Please indicate date(s) of episode(s).				
3. Is your client on any medications?				
☐ Yes. Please give details.				
□No				
4. Does your client have a history of substance	e abuse (alcohol or drugs)?			
☐ Yes. Please give details				
□No				
5. Has your client been hospitalized, required symptoms or treatment?	ECT, been seen in the eme	rgency room, or been on disability for psychiatric		
☐ Yes. Please give details				
□No				
6. Has your client smoked cigarettes in the la	st 12 months?			
Yes				
□No				
7. Does your client have any other major healt	th problems (e.g., cancer, e	tc.)?		
☐ Yes. Please give details				
□ No				